

Pest Management Methods



PENN STATE



College of
Agricultural Sciences
Cooperative Extension

Penn State College of Agricultural Sciences research, extension, and resident education programs are funded in part by Pennsylvania counties, the Commonwealth of Pennsylvania, and the U.S. Department of Agriculture.

This publication is available from the Publications Distribution Center, The Pennsylvania State University, 112 Agricultural Administration Building, University Park, PA 16802. For information telephone (814) 865-6713.

Where trade names appear, no discrimination is intended, and no endorsement by the Penn State College of Agricultural Sciences is implied.

Issued in furtherance of Cooperative Extension Work, Acts of Congress May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture and the Pennsylvania Legislature. T. R. Alter, Director of Cooperative Extension, The Pennsylvania State University.

This publication is available in alternative media on request.

The Pennsylvania State University is committed to the policy that all persons shall have equal access to programs, facilities, admission, and employment without regard to personal characteristics not related to ability, performance, or qualifications as determined by University policy or by state or federal authorities. The Pennsylvania State University does not discriminate against any person because of age, ancestry, color, disability or handicap, national origin, race, religious creed, sex, sexual orientation, or veteran status. Direct all inquiries regarding the nondiscrimination policy to the Affirmative Action Director, The Pennsylvania State University, 201 Willard Building, University Park, PA 16802-2801; Tel. (814) 865-4700/V; (814) 863-1150/TTY.

© The Pennsylvania State University 1998

5M998NVO

People who practice IPM (integrated pest management) understand that eradicating insect pests and diseases of plants is usually unrealistic. IPM primarily consists of methods used to prevent plant problems from occurring in the first place. To practice IPM in the landscape, choose plants that are well suited to the site. Plant them properly and keep the plants healthy by carefully watering, fertilizing, and pruning them. Watch out for problems on your plants as they arise. If a pest or disease causes unacceptable damage despite preventative efforts, choose an effective management method that will have the least amount of impact on other living creatures and the environment.

FOLLOW THESE STEPS TO MAKE AN IPM CONTROL DECISION:

1. CONFIRM THAT THERE IS A PEST PROBLEM

Look for pests and diseases and the evidence or signs they leave. Look for symptoms the plant exhibits as a result of pest activity. Examine your plants often. Identify your plants to be sure that the twisted leaves, unusual coloration, or strange-looking structures you see are not a normal part of the plant. Try to rule out site-related problems by making sure that the soil type, drainage conditions, fertility level, and other environmental conditions are favorable for the plant.

2. IDENTIFY THE PROBLEM

Effective pest management depends on the accurate identification of the pest. Insects and mites often are associated with specific plants, and they follow certain development and behavior patterns as the season progresses. Use reference books from the library or garden center to identify pests. If you can't find an accurate description there, consult someone in your local extension office. Learn about the insect's life cycle, behavior, and natural enemies.

Plant diseases may be caused by pathogens including fungi, nematodes, bacteria, or viruses. Each pathogen is capable of infecting only certain plants. Infection occurs under particular environmental conditions, with symptoms of

the disease appearing later. To identify plant diseases accurately, compare visible signs and symptoms of the disease with descriptions in reference books. Some diseases are more difficult to identify, and you may need a laboratory analysis or the help of an expert. These services are available through landscape professionals or your local extension office.

3. DETERMINE IF A CONTROL MEASURE IS NEEDED

Determine if the damage is severe enough to justify a management tactic. Is the damage actually affecting the health of the plant? If not, does it make the plant look bad enough to detract from the appearance of your landscape? You may become alarmed if you notice that caterpillars are making holes in leaves, but if the damage is slight or occurs late in the season, you probably won't need to control the caterpillars.

Are the pest's natural enemies present? In most cases, they will be. If so, an application of a nonselective insecticide could kill them, allowing the pest population to rebound uninhibited by predators and parasites, which may have been providing significant control.

4. CHOOSE A METHOD

If a control is needed, consider physical or biorational methods first. If they are unavailable or impractical, you may need to carefully use a conventional chemical control.

Physical methods

Pests can be removed from plants physically. For example, some aphids and mites can be knocked off by spraying the plant with water. Bagworm larvae can be picked off an infested plant.

You can use traps to catch certain pests, and barriers to protect plants from insect attack or disease infection. One effective method for controlling gypsy moth larvae on small numbers of trees is to put a band of folded burlap around the tree trunk to provide an artificial resting site for the caterpillars, and then destroy the caterpillars that gather there. Applying an antitranspirant spray to lilac leaves in summer to prevent infection by the spores of powdery mildew is another example of a protective barrier.

In some cases, the best solution may be physically removing the plant and

replacing it with one that will not be affected by the pest or disease. Thinning crowded plants to improve air circulation can reduce many disease problems.

Biorational methods

Biorational methods can be divided into two groups. The first group includes living organisms that can kill the pest. The second group includes naturally occurring biochemicals that are harmful to the pest yet often are harmless to other living organisms.

Insect pests frequently have natural enemies that are beneficial to the landscape. These beneficial insects often exist in the landscape naturally, but they also can be introduced. "Beneficials" may be predators or parasites. One common example of a beneficial predator is the lady beetle. Both the larvae and adult lady beetles eat aphids and other soft-bodied insects. Other predators include lacewings, spined soldier bugs, flower flies, and spiders. Parasites live on and often kill another organism, called the host. Some parasitic wasps use caterpillars, whiteflies, aphids, and soft scales as hosts.

An example of a method that uses a naturally occurring biochemical is the bacterium *Bacillus thuringiensis* (Bt). Bt contains a protein that is poisonous to specific insects, yet harmless to other organisms. Bt can be sprayed on plants. When the sensitive insect pest feeds on the sprayed leaves, it will ingest the protein and be killed.

Chemicals

Conventional chemicals are used only as a last resort in an IPM program, but sometimes are the most effective means of control. To have the greatest effect, these materials need to be applied on a specific part of the plant when the pest is most vulnerable. Always apply chemical controls according to label directions.

In many cases, environmentally safe pesticides such as horticultural oil or insecticidal soap are effective choices. Again, applications must be timed carefully to have the greatest effect on the pest insect population. Because they have no residual activity after they have dried, soaps and oils are usually the option that is the least disruptive to populations of beneficial organisms.

The label of every pesticide formula-tion displays a signal word that relates

to the pesticide's level of toxicity. These words, from least toxic to most toxic are: "caution," "warning," and "danger." Use these words as guidelines to help you choose the least hazardous material among the effective alternatives. For most landscape pests, you need to consider pesticides in only the first two categories. Some pesticide formulations can be applied only by applicators with special training and who are certified by the state's department of agriculture.

A pesticide may be applied only to plants that are listed on the label. Refer to the *Woody Ornamental Insect, Mite, and Disease Management Guide*, published by Penn State Cooperative Extension, or to another current reference for a list of materials that are registered for use on plants in your state. Read the label to be sure that you have chosen a product that is effective against the pest you have identified, and choose your timing based on professional recommendations.

Follow these procedures for your safety and success:

- Mix pesticides according to label instructions. Do not use more or less concentrate in the mixture than the label recommends. Mix only as much material as you need for the application.
- Wear protective clothing as specified on the label.
- Label a set of mixing and measuring tools that are used *only* for insecticides and fungicides, and store them with the products. Use a separate set of measuring tools and spray equipment for herbicides.
- Keep pets and people away from the area where you store, mix, and apply pesticides. Stay away from a treated area for as long as the label directs.
- Do not spray on a windy day or when air temperatures will be above 85°F before the spray solution dries.
- Clean equipment and mixing tools as soon as you finish spraying.
- After spraying, change your protective clothing and bathe. Wash the clothes you were wearing separately from your regular laundry.
- Keep records of where and when you sprayed, what pesticide you used, and how much you used. Give the treatment time to work, then evaluate and record your results.

REFERENCE

Woody Ornamental Insect, Mite, and Disease Management Guide. 1995. Penn State Cooperative Extension. Penn State Publications Distribution Center, 112 Agricultural Administration Building, University Park, PA 16802-2600.

FOR MORE INFORMATION

Penn State Cooperative Extension, Delaware Cooperative Extension, and the Southeast Pennsylvania IPM Research Group have been working together to provide information and educational materials on IPM and landscaping.

This fact sheet, "Pest Management Methods," is part of a series of educational fact sheets about understanding and using Integrated Pest Management. Other topics in the series include:

- Creating Healthy Landscapes—Introduction
- Choose Plants Wisely
- Plant With Care
- Promote Plant Health
- Keep Plants Well Groomed
- Monitor Pests and Keep Records

Copies are available from your local extension office.

The Southeast Pennsylvania IPM Research Group is a collaboration of university and industry horticulture professionals who are inspecting landscapes across the region to monitor pest populations and share current IPM data. The group is partially supported by the Pennsylvania IPM Program (PAIPM). For more information about the research group, contact: Penn State Cooperative Extension, Montgomery County, 1015 Bridge Road, Suite H, Collegeville, PA 19426-1179; telephone: (610) 489-4315.

Prepared by:
Stevie Daniels, Penn State master gardener
Gregory Hoover, extension entomologist
Larry Kuhns, professor of horticulture
Gary Moorman, professor of plant pathology
Robert Nuss, professor of horticulture
David Suchanic, Penn State Cooperative Extension agent
Emelie Swackhamer, Penn State Cooperative Extension agent
Title illustration by Mary Thompson